

UDC 593.12(265.54)

***CEPHALOGULLMIA CAUDATA* GEN. NOV., SP. NOV. (RHIZARIA, FORAMINIFERA),  
A MONOTHALAMOUS FORAMINIFERA FROM THE SEA OF JAPAN**

© 2025 **N. Sergeeva** and **O. Anikeeva**

A. O. Kovalevsky Institute of Biology of the Southern Seas of RAS, Sevastopol, Russian Federation

E-mail: [alegria@ibss-ras.ru](mailto:alegria@ibss-ras.ru)

Received by the Editor 02.07.2024; after reviewing 18.10.2024;  
accepted for publication 20.03.2025.

We describe a new species and genus of soft-walled monothalamous foraminifera from the Sea of Japan. The material was sampled off the Primorsky shelf during the 64<sup>th</sup> cruise of the RV “Akademik Oparin” (17 June to 08 July, 2021). Bottom sediments were sampled at 17 stations in a depth range of 0.3–86 m. The species described here was found at depths of 13 and 86 m. An elongated organic-walled test of *Cephalogullmia caudata* sp. nov. has an inflated anterior (apertural) part which gradually narrows towards the adapertural end. The length varies 600 to 1,180 µm; the width at the widest part (the ‘head’) is 50 to 135 µm, and at the narrowest part, it is 10 to 20 µm. A single apertural structure (10–28 µm in diameter) is located at the widest end of the test. The apertural structure is slightly elevated; in some specimens, it resembles a nipple. The nucleus is visible only in some specimens; its size is from 16 × 12 µm to 30 × 15 µm. The comparative analysis shows that the new genus *Cephalogullmia* differs from *Micrometula* Nyholm, 1952 in the shape of the test: it has more rounded and inflated apertural end, compared to that of the slender conical test of *Micrometula*. The new genus differs from *Cylindrogullmia* Nyholm, 1974 in the shape of the test as well: in the latter one, it is cylindrical. It also differs from another organic-walled monothalamid with the elongated test, *Bowseria* Sinniger et al., 2008, in having a darker protoplasm and a narrowed (tail-shaped) adapertural end.

**Keywords:** soft-walled foraminifera, Sea of Japan, *Cephalogullmia caudata* gen. nov., sp. nov., monothalamids

Soft-walled monothalamous foraminifera are an important component of meiofauna in many coastal and deepwater habitats [Gooday, 2002; Majewski et al., 2007; Sergeeva, Anikeeva, 2018, 2023], as well as in freshwater ones [Holzmann et al., 2021]. In recent years, scientific interest in this group has increased significantly [Anikeeva et al., 2013; Sergeeva et al., 2010]. Although molecular analysis is now available for many monothalamous foraminifera [Gooday et al., 2011, 2022; Lejzerowicz et al., 2015; Siemensma et al., 2021; Voltski, Pawłowski, 2015], some recent taxonomic descriptions still rely on morphological characteristics alone [Anikeeva et al., 2013; Golemansky, 1999; Gooday et al., 2006; Henderson, 2023; Ohkawara et al., 2009; Sergeeva, Anikeeva, 2008, 2020, 2021]. Here, we describe a new monothalamous taxon, an organic-walled ‘allogromiid,’ based on its morphological features.

Studies of meiobenthos in the Sea of Japan have been mainly focused on multicellular eukaryotes, especially the dominant group, free-living nematodes. Out of unicellular eukaryotes, only hard-shelled foraminifera have been considered as a component of meiobenthic communities [Preobrazhenskaya, Tarasova, 1990]. The results of detailed studies of the species diversity of hard-shelled foraminifera in the Sea of Japan are provided by T. Lukina and T. Tarasova [2013] in their check-list of free-living

invertebrates in Far Eastern seas of Russia – a publication covering 217 foraminifera species. A representative of a genus of soft-walled foraminifera, *Allogromia* sp. (the family Allogromiidae), is reported to be found in the Bering Sea and off the South Kuril Islands and northeastern Sakhalin Peninsula. This appears to be the only record of soft-walled foraminifera in the Northwestern Pacific reflecting the lack of information on these protists, their specific morphological characteristics, and the need to apply special methods to study them.

For the first time, soft-walled foraminifera were investigated in Far Eastern waters of Russia (the Sea of Japan) in summer 2021, during the research cruise of the RV “Akademik Oparin.” In total, 49 species and morphotypes of monothalamids representing the families Allogromiidae and Saccamminidae were recorded in this area [Sergeeva, Anikeeva, 2023]. The fauna of monothalamids in the Sea of Japan included the one that we describe here as the type species of a new genus.

The aim of this paper is to provide a description of *Cephalogullmia caudata* gen. nov., sp. nov., an inhabitant of the Sea of Japan (the Northwestern Pacific).

## MATERIAL AND METHODS

Material was sampled off the Primorsky shelf, the Sea of Japan, during the 64<sup>th</sup> cruise of the RV “Akademik Oparin” (17 June to 08 July, 2021). Bottom sediments were sampled at 17 stations in a depth range of 0.3 to 86 m. The species described was found at sta. 1 and 40 (depths of 13 and 86 m). At shallower stations, divers sampled sediment cores (two replicates), 5 cm long, with a cross-sectional area of 10 cm<sup>2</sup>. At greater depths, similar cores were taken from sediment obtained with a Van Veen grab sampler. Samples of bottom sediments were fixed onboard the RV with 75% ethanol and later processed and analyzed in IBSS laboratory (Sevastopol, Russian Federation). Samples were washed through two sieves: an upper one with a mesh size of 1 mm and a lower one with a mesh size of 63 µm. For the second series of samples, we used a lower sieve with a mesh size of 32 µm.

Detailed microscopic analysis of soft-walled foraminifera was performed under MSP-2, Olympus CX41, Nikon, and Mikmed-6 microscopes equipped with photo cameras.

## RESULTS

**Taxonomy.** According to traditional morphology-based classification systems [Kaminski, 2004; Loeblisch, Tappan, 1988], the new single-chambered (monothalamous) taxa described here belong to the sub-order Allogromiina (organic-walled). However, recent molecular phylogenetic studies, which revealed a series of monophyletic clades that cut across morphology-based taxa, have undermined the traditional higher-level classification of monothalamous organic-walled and agglutinated foraminifera [Habura et al., 2008; Pawlowski et al., 2003, 2013]. Therefore, we avoid formal higher taxa and place the new genera in the informal group of ‘monothalamids’ [Pawlowski et al., 2013].

Rhizaria Cavalier-Smith, 2002

Foraminifera d’Orbigny, 1826

‘Monothalamids’ Pawlowski et al., 2013

Genus *Cephalogullmia* Sergeeva et Anikeeva, gen. nov.


Type species *Cephalogullmia caudata* Sergeeva et Anikeeva, sp. nov.

**Diagnosis of the genus.** A monothalamous foraminifera having an elongated organic-walled test with an inflated anterior (apertural) part that gradually narrows towards the adapertural end (Fig. 1). The width at the inflated front part (the ‘head’) exceeds the width of the body by 5–6 times at the narrowest

part. A single slightly elevated apertural structure is located at the widest end of the test. In some specimens, it resembles a nipple. The oval nucleus is localized in the widened part of the body before its tapering and is visible only in some specimens.

**Derivation of the genus name.** The name of the genus indicates its similarity to the genus *Nemogullmia* in terms of test length and the rounded apertural end resembling a head (in Latin, ‘cephalo’).

**Remarks.** Comparative analysis showed that the investigated specimens cannot be related to any of the known foraminifera genera. We made a comparative diagnosis of the new genus with three morphologically most similar genera of soft-walled foraminifera. The key difference between the new genus *Cephalogullmia* and *Micrometula* Nyholm, 1952 is in the shape of the test: there is the test with a more rounded and inflated apertural end than the slender conical test of *Micrometula*. The new genus differs from *Cylindrogullmia* Nyholm, 1974 by the presence of a sharply tapering ‘tail’ part, while the test of *Cylindrogullmia* specimens has approximately the same diameter throughout the body length. It also differs from another elongated organic-walled monothalamid, *Bowseria* Sinniger et al., 2008, in having a darker protoplasm and a narrowed (tail-shaped) adapertural end.



**Fig. 1.** Schematic image of *Cephalogullmia caudata*: 1, aperture; 2, organic test wall; 3, nucleus; 4, cytoplasm; 5, ‘tail’ part

*Cephalogullmia caudata* Sergeeva et Anikeeva, sp. nov. (Fig. 2).

Syn. Allogromiid sp. 9J [Sergeeva, Anikeeva, 2023, pp. 446–447, Fig. 9e, f].

**Diagnosis of species.** The elongated organic-walled test has an inflated anterior (apertural) part which gradually narrows towards the adapertural end. The body length varies from 600 to 1,180  $\mu\text{m}$ ; the width at the widest part (the ‘head’) is 50 to 135  $\mu\text{m}$ , and at the narrowest part, it is 10 to 20  $\mu\text{m}$ . A single apertural structure (10–28  $\mu\text{m}$  in diameter) is located at the widest end of the test. The apertural structure is slightly elevated; in some specimens, it resembles a nipple. The nucleus is visible only in some specimens, and its size is from  $16 \times 12 \mu\text{m}$  to  $30 \times 15 \mu\text{m}$ . Coefficient C (length-to-width ratio) ranges 8.7 to 16.5 (based on the maximum width of the body).

**Derivation of specific name.** Latin ‘caudata’ = ‘tail’ refers to the shape of the main part of the body.

**Type material.** The type specimen (registration number Meib.93.Al.t) and two paratype specimens (registration numbers Meib.94.Al.p.1 and Meib.95.Al.p.2) are mounted in water–glycerol preparations and stored at IBSS. Previous experience has shown that similar organic-walled specimens are too delicate to be mounted in either Canada balsam or a gelatin–glycerol mixture.

**Type locality.** The type specimen and the paratype specimens (Figs 2–4) are from the Sea of Japan ( $43^{\circ}42.7'N$ ,  $135^{\circ}25.6'E$ ; depth of 86 m). They were picked from samples taken during the research cruise of the RV “Akademik Oparin,” 17 June to 08 July, 2021.

**Other material.** Three more specimens from the type locality (Fig. 5) were examined.



**Description.** The type specimen has an elongated shape (Fig. 2A). The test is organic, thin, and transparent. It gradually narrows from the apertural end to the adapertural one. The cytoplasm is homogeneous and fine-grained. The single aperture is located at the widest end of the test and is 16  $\mu\text{m}$  in diameter (Fig. 2B). The test length is 1,070  $\mu\text{m}$ , and the maximum width is 90  $\mu\text{m}$ . The nucleus is ellipsoidal, oblong (Fig. 2C); it is located in the middle part of the cell, and it is  $30 \times 15 \mu\text{m}$ . Coefficient C is about 11.8 (based on the maximum width of the body).




**Fig. 2.** The type specimen: A, general view; B, aperture; C, nucleus. Scale bars are 100  $\mu\text{m}$  (A) and 20  $\mu\text{m}$  (B, C)

The paratype specimen 1 (Fig. 3A) is 1,150  $\mu\text{m}$  in length and tapers from 110  $\mu\text{m}$  in width at the widest point to 20  $\mu\text{m}$  at the narrow adapertural end (Fig. 3B). The aperture has a diameter of 28  $\mu\text{m}$ . Coefficient C is 10.4.




**Fig. 3.** The paratype specimen 1: A, general view; B, aperture. Scale bars are 200  $\mu\text{m}$  (A) and 20  $\mu\text{m}$  (B)

The paratype specimen 2 (Fig. 4A) is 1,180  $\mu\text{m}$  in length and tapers from 135  $\mu\text{m}$  in width at the widest point to 25  $\mu\text{m}$  at the narrow adapertural end. The aperture (Fig. 4B) is located at the wider end of the body; however, its size cannot be measured, as it is not clearly visible. Coefficient C is 8.7. The nucleus is not clearly visible in either of the paratypes.



**Fig. 4.** The paratype specimen 2: A, general view; B, aperture. Scale bars are 100  $\mu\text{m}$  (A) and 50  $\mu\text{m}$  (B)





**Fig. 5.** Specimen 1J ('J' indicates the initial letter in the name of the Sea of Japan): A, general view; B, aperture. Specimen 2J: C, general view; D, aperture. Specimen 3J: E, general view; F, aperture; G, nucleus. Scale bars are 100 µm (A, C, E) and 20 µm (B, D, F, G). Specimens 1J and 2J are partially deformed

We examined in detail only 6 individuals (Table 1), because very delicate and fragile tests of other specimens were significantly deformed or damaged during the samples washing.

**Table 1.** Morphological parameters (in µm) of the type specimen, the two paratype specimens, and other examined specimens of *Cephalogullmia caudata* (–, not visible)

Specimen	Test length	Test width (maximum)	Diameter of aperture	Nucleus	Coefficient C
Type specimen	1,070	90	16	30 × 15	11.8
Paratype specimen 1	1,150	110	28	–	10.4
Paratype specimen 2	1,180	135	–	–	8.7
Specimen 1J	600	50	13	–	12
Specimen 2J	1,160	70	26	–	16.5
Specimen 3J	640	50	10	16 × 12	12.8

This work was carried out within the framework of IBSS state research assignment “Biodiversity as the basis for the sustainable functioning of marine ecosystems, criteria and scientific principles for its conservation” (No. 124022400148-4).

**Acknowledgement.** We express our deep gratitude to the administration of the Pacific Institute of Bioorganic Chemistry, Far Eastern Branch of the Russian Academy of Sciences, for the opportunity to sample bottom sediments in the Sea of Japan, as well as to the captain of the RV “Akademik Oparin” and our colleagues S. Trofimov and Yu. Litvin (department of benthic ecology, IBSS) who provided invaluable assistance in obtaining scientific material. We are very grateful to Prof. A. Gooday for his helpful consultations and revision of the English text of the paper. We thank the anonymous reviewers who provided a number of useful tips and recommendations for improving our manuscript.

## REFERENCES

1. Anikeeva O. V., Sergeeva N. G., Gooday A. J. Two new genera and species of the monothalamous foraminifera from coastal waters of the Black Sea. *Marine Biodiversity*, 2013, vol. 43, iss. 4, pp. 473–479. <https://doi.org/10.1007/s12526-013-0177-0>
2. Golemansky V. G. *Lagynis pontica* n. sp., a new monothalamous rhizopod (Granuloreticulosea: Lagynidae) from the Black Sea littoral. *Acta Zoologica Bulgarica*, 1999, vol. 51, pp. 3–13.
3. Gooday A. J. Organic-walled allogromiids: Aspects of their occurrence, diversity and ecology in marine habitats. *Journal of Foraminiferal Research*, 2002, vol. 32, no. 4, pp. 384–399. <https://doi.org/10.2113/0320384>
4. Gooday A. J., Anikeeva O. V., Sergeeva N. G. *Tinogullmia lukyanovae* sp. nov. – a monothalamous, organic-walled foraminiferan from the coastal Black Sea. *Journal of the Marine Biological Association of the United Kingdom*, 2006, vol. 86, iss. 1, pp. 43–49. <https://doi.org/10.1017/S0025315406012847>
5. Gooday A. J., Anikeeva O. V., Pawłowski J. New genera and species of monothalamous foraminifera from Balaclava and Kazach'ya Bays (Crimean Peninsula, Black Sea). *Marine Biodiversity*, 2011, vol. 41, iss. 4, pp. 481–494. <https://doi.org/10.1007/s12526-010-0075-7>
6. Gooday A. J., Holzmann M., Schwarzgruber E., Cedhagen T., Pawłowski J. Morphological and molecular diversity of monothalamids (Rhizaria, Foraminifera), including two new species and a new genus, from SW Greenland. *European Journal of Protistology*, 2022, vol. 86, art. no. 125932 (25 p.). <https://doi.org/10.1016/j.ejop.2022.125932>
7. Habura A., Goldstein S. T., Broderick S., Bowser S. S. A bush, not a tree: The extraordinary diversity of cold-water basal foraminiferans extends to warm-water environments. *Limnology and Oceanography*, 2008, vol. 53, iss. 4, pp. 1339–1351. <https://doi.org/10.4319/lo.2008.53.4.1339>
8. Henderson Z. Soft-walled monothalamid foraminifera from the intertidal zones of the Lorn area, north-west Scotland. *Journal of the Marine Biological Association of the United Kingdom*, 2023, vol. 103, art. no. e18 (21 p.). <https://doi.org/10.1017/S0025315423000061>
9. Holzmann M., Gooday A. J., Siemensma F., Pawłowski J. Review: Freshwater and soil foraminifera – a story of long-forgotten relatives. *Journal of Foraminiferal Research*, 2021, vol. 51, no. 4, pp. 318–331. <https://doi.org/10.2113/gsjfr.51.4.318>
10. Kaminski M. A. The year 2000 classification of the agglutinated foraminifera. In: *Proceedings of the Sixth International Workshop on Agglutinated Foraminifera, Prague, Czech Republic, 1–7 September, 2001* / M. Bubik, M. A. Kaminski (Eds). Kraków, Poland : Grzybowski Foundation, 2004, pp. 237–255. (Grzybowski Foundation, Special Publication ; no. 8).
11. Lejzerowicz F., Voltski I., Pawłowski J.



- Foraminifera of the Kuril–Kamchatka Trench area: The prospects of molecular study. *Deep Sea Research Part II: Topical Studies in Oceanography*, 2015, vol. 111, pp. 19–25. <https://doi.org/10.1016/j.dsr2.2014.10.003>
12. Loeblich A. R., Tappan J. H. *Foraminiferal Genera and Their Classification*. New York : Van Nostrand Reinhold Company, 1988, 970 p. <https://doi.org/10.1007/978-1-4899-5760-3>
  13. Lukina T. G., Tarasova T. S. Phylum Sarcomastigophora, class Granuloreticulosa. In: *Check-List of Species of Free-Living Invertebrates of the Russian Far Eastern Seas* / B. I. Sirenko (Ed.). Saint Petersburg : [ZIN RAS], 2013, pp. 16–24. (Explorations of the Fauna of the Seas ; 75 (83)). (in Russ.)
  14. Majewski W., Lecroq B., Sinniger F., Pawłowski J. Monothalamous foraminifera from Admiralty Bay, King George Island, West Antarctica. *Polish Polar Research*, 2007, vol. 28, no. 3, pp. 187–210.
  15. Ohkawara N., Kitazato H., Uematsu K., Gooday A. J. A minute new species of *Saccamina* (monothalamous foraminifera; Protista) from the abyssal Pacific. *Journal of Micropalaeontology*, 2009, vol. 28, iss. 2, pp. 143–151. <https://doi.org/10.1144/jm.28.2.143>
  16. Pawłowski J., Holzmann M., Berney F. C., Gooday A. J., Cedhagen T., Habura A., Bowser S. S. The evolution of early Foraminifera. *Proceedings of the National Academy of Sciences*, 2003, vol. 100, iss. 20, pp. 11494–11498. <https://doi.org/10.1073/pnas.2035132100>
  17. Pawłowski J., Holzmann M., Tyszká J. New supraordinal classification of Foraminifera: Molecules meet morphology. *Marine Micropalaeontology*, 2013, vol. 100, pp. 1–10. <https://doi.org/10.1016/j.marmicro.2013.04.002>
  18. Preobrazhenskaya T. V., Tarasova T. S. Donnye foraminifery nekotorykh raionov zaliva Petra Velikogo. In: *Rasprostranenie i ekologiya sovremennykh i iskopaemykh morskikh organizmov*. Vladivostok : Dal'nevostochnoe otdelenie Akademii nauk SSSR, 1990, pp. 11–18. (in Russ.)
  19. Sergeeva N. G., Anikeeva O. V. *Goodayia rostellatum* gen. n., sp. n. (Protozoa) – a monothalamous foraminifera from the Black Sea. *Vestnik zoologii*, 2008, vol. 42, no. 5, pp. 467–471. <https://elibrary.ru/jxfqsd>
  20. Sergeeva N. G., Anikeeva O. V. *Soft-shelled Foraminifera of the Black Sea and the Sea of Azov*. Simferopol : ARIAL, 2018, 156 p. (in Russ.). <https://doi.org/10.21072/978-5-907118-84-3>
  21. Sergeeva N. G., Anikeeva O. V. New Black Sea monothalamous Foraminifera from the genus *Nemogullmia* Nyholm, 1953 (Allogromiida: Shepheardellinae). *Invertebrate Zoology*, 2020, vol. 17, no. 2, pp. 176–188. <https://doi.org/10.15298/invertzool.17.2.07>
  22. Sergeeva N. G., Anikeeva O. V. *Vellaria solenta* (Monothalamaea: Allogromiidae) – new species of soft-walled foraminifera from Sivash Bay (the Sea of Azov). *Invertebrate Zoology*, 2021, vol. 18, no. 2, pp. 152–158. <https://doi.org/10.15298/invertzool.18.2.06>
  23. Sergeeva N. G., Anikeeva O. V. First investigations of benthic soft-walled foraminifera and gromiids (Protozoa) in the northwestern Sea of Japan. *Russian Journal of Marine Biology*, 2023, vol. 49, no. 6, pp. 435–452. <https://doi.org/10.1134/S106307402306007X>
  24. Sergeeva N. G., Anikeeva O. V., Gooday A. J. Soft-shelled, monothalamous foraminifera from the oxic/anoxic interface (NW Black Sea). *Journal of Micropalaeontology*, 2010, vol. 56, no. 3–4, pp. 393–407. <https://elibrary.ru/xmdoph>
  25. Siemensma F., Holzmann M., Apothéloz-Perret-Gentil L., Clauß S., Voelcker E., Bettighofer W., Khanipour Roshan S., Walden S., Dumack K., Pawłowski J. Broad sampling of monothalamids (Rhizaria, Foraminifera) gives further insight into diversity of non-marine foraminifera. *European Journal of Protistology*, 2021, vol. 77, art. no. 125744 (21 p.). <https://doi.org/10.1016/j.ejop.2020.125744>
  26. Voltski I., Pawłowski J. *Flexammina islandica* gen. nov., sp. nov. and some new phylotypes of monothalamous foraminifera from the coast of Iceland. *Zootaxa*, 2015, vol. 3964, no. 2, pp. 245–259. <https://doi.org/10.11646/zootaxa.3964.2.5>

***CEPHALOGULLMIA CAUDATA* GEN. NOV., SP. NOV. (RHIZARIA, FORAMINIFERA) —  
МОНОТАЛАМУСНАЯ ФОРАМИНИФЕРА ИЗ ЯПОНСКОГО МОРЯ**

**Н. Г. Сергеева, О. В. Аникеева**

ФГБУН ФИЦ «Институт биологии южных морей имени А. О. Ковалевского РАН»,  
Севастополь, Российская Федерация  
E-mail: [alegria@ibss-ras.ru](mailto:alegria@ibss-ras.ru)

Мы приводим описание нового вида и рода мягкораковинных моноталамусных фораминифер из Японского моря. Материал собран в районе приморского шельфа во время 64-го рейса НИС «Академик Опарин» (17 июня — 08 июля 2021 г.). Пробы донных осадков отобраны на 17 станциях в диапазоне глубин от 0,3 до 86 м. Описываемый здесь вид был найден на глубинах 13 и 86 м. Удлиненная органическая стенка раковины *Cephalogullmia caudata* sp. nov. имеет раздутую переднюю (апертурную) часть, которая постепенно сужается к дистальному концу. Длина тела варьирует от 600 до 1180 мкм; ширина в самой широкой части («голове») — от 50 до 135 мкм, в самой узкой части — от 10 до 20 мкм. Единственная апертурная структура (диаметр 10–28 мкм) расположена на самом широком конце раковины. Апертура слегка приподнята, у некоторых экземпляров напоминает сосок. Ядро видно только у некоторых экземпляров; его размеры составляют от 16 × 12 мкм до 30 × 15 мкм. Сравнительный анализ показал, что новый род *Cephalogullmia* отличается от *Micrometula* Nyholm, 1952 по форме раковины: она более округлая и раздутая в апертурной части, чем узкая раковина конической формы у *Micrometula*. От *Cylindrogullmia* Nyholm, 1974 новый род также отличается по форме раковины: у *Cylindrogullmia* она больше напоминает цилиндр. Описываемый нами род отличается и от другой моноталамиды с удлиненной раковинной и органическими стенками — *Bowseria* Sinniger et al., 2008 — более темной протоплазмой и суженным (хвостобразным) дистальным концом.

**Ключевые слова:** мягкораковинные фораминиферы, Японское море, *Cephalogullmia caudata* gen. nov., sp. nov., моноталамиды